

The Evening Sky Map

FREE* EACH MONTH FOR YOU TO EXPLORE, LEARN & ENJOY THE NIGHT SKY

WWW.SKYPMAPS.COM

NORTHERN HEMISPHERE
JANUARY 2006

Sky Calendar – January 2006


- 1 **Moon near Venus** at 13h UT (19° from Sun, evening sky).
- 1 **Moon at perigee** (closest to Earth) at 23h UT (distance 361,750 km; size 33.0').
- 3 **Quadrantid Meteor Shower** peaks at 16:20 UT. Produces medium-speed meteors (up to 100 per hour). Radiant in the constellation Boötes. Parent comet is unknown.
- 4 **Earth at Perihelion** (closest to Sun) at 16h UT. The Sun-Earth distance is 0.983327 a.u. or about 147.1 million kilometers.
- 6 **First Quarter Moon** at 18:56 UT.
- 8 **Moon near Mars** at 19h UT (evening sky). Mag. -0.4.
- 10 **Moon very near the Pleiades** at 2h UT (evening sky). Occultation visible from North America and northern parts of South America.
- 13 **Venus at inferior conjunction** with the Sun at 24h UT. The planet passes into the morning sky.
- 14 **Moon near Pollux** at 8h UT (midnight sky).
- 14 **Full Moon** at 9:48 UT. The full Moon of January is called the "Old Moon" or the "Moon After Yule" in North American folklore.
- 15 **Moon near Saturn** at 16h UT (morning sky). Mag. -0.2.
- 17 **Moon at apogee** (farthest from Earth) at 19h UT (distance 405,885 km; angular size 29.4').
- 17 **Moon near Regulus** at 10h UT (morning sky).
- 21 **Moon very near Spica** at 21h UT (morning sky). Occultation visible from Asia and Indonesia.
- 22 **Last Quarter Moon** at 15:14 UT.
- 23 **Moon near Jupiter** at 17h UT (morning sky). Mag. -1.9.
- 25 **Moon very near Antares** at 10h UT (morning sky). Occultation visible from southern Mexico, Peru, and Ecuador.
- 27 **Saturn at opposition** (opposite the Sun) at 23h UT. Visible all night long, the ringed planet is at its brightest (mag. -0.2) and closest (disk diameter 20.5") all year. Saturn's rings appear magnificent even in a small telescope.
- 29 **New Moon** at 14:15 UT. Beginning of lunation 1028.
- 30 **Moon at perigee** (closest to Earth) at 8h UT (distance 357,778 km; size 33.4').

All times in Universal Time (UT). (USA Eastern Standard Time = UT - 5 hours.)



SAVE ON RECOMMENDED PRODUCTS • <http://Skymaps.com/store>

- STAR ATLAS & PLANISPHERES
- DESKTOP PLANETARIUM SOFTWARE
- LARGE-FORMAT HUBBLE PRINTS
- STAR CHARTS & ASTRO POSTERS
- TELESCOPES FOR BEGINNERS
- 2006 ASTRONOMY CALENDARS



SKY MAP SHOWS HOW THE NIGHT SKY LOOKS

EARLY JAN 8 PM
LATE JAN 7 PM

SKY MAP DRAWN FOR A LATITUDE OF 40° NORTH AND IS SUITABLE FOR LATITUDES UP TO 15° NORTH OR SOUTH OF THIS

From northern latitudes, stars appear to rotate around the North Celestial Pole (NCP). Compass directions are indicated along the horizon circle. (For example, "North", "South", "East", and "West" are the directions you would see if you were standing on the Earth's surface.)

The Milky Way (the band of light) is our home galaxy. Dark stars are required to see it.

CELESTIAL OBJECTS ARE LOCATED BETWEEN THE ZENITH AND THE HORIZON.

THE CENTER OF THE MAP IS THE PART OF THE SKY DIRECTLY OVERHEAD (ZENITH) AND THE OUTER CIRCLE IS THE HORIZON. (The map is drawn for a latitude of 40° north.)

Can you see the Andromeda Galaxy (M31) with your eyes? It is a galaxy 2.93 million light-years away!

The Pleiades, or Seven Sisters, are easily seen with the naked eye. The V-shaped Hyades are nearby.

INSTRUCTIONS: THE SKY MAP SHOWS THE ENTIRE NIGHT SKY FROM HORIZON-TO-HORIZON AS IT APPEARS ON CERTAIN DATES AND TIMES.

THE THREE BRIGHTEST STARS "in a line" in the southeast form what is known as the "Belt of Orion".

THE COMPASS DIRECTION THAT APPEARS ALONG THE BOTTOM OF THE MAP IS THE SAME AS THE DIRECTION THAT YOU FACE. BEGIN BY USING THE SKY MAP TO FIND A BRIGHT STAR PATTERN IN THE SKY.

Symbols

- Galaxy ☾
- Double Star ●
- Variable Star ⊙
- Diffuse Nebula □
- Planetary Nebula ◇
- Open Star Cluster ○
- Global Star Cluster ⊕

Star Magnitudes ● -1 0 1 2 3 4

Copyright © 2000–2006 Kym Thalassoudis. All Rights Reserved.

* TERMS OF USE: FREE FOR NON-COMMERCIAL EDUCATIONAL USE. ASTRONOMY EDUCATION GROUPS MAY FREELY DISTRIBUTE PRINTED HANDOUTS. FULL DETAILS AT <http://Skymaps.com/terms.html>

About the Celestial Objects

Listed on this page are several of the brighter, more interesting celestial objects visible in the evening sky this month (refer to the monthly sky map). The objects are grouped into three categories. Those that can be easily seen with the naked eye (that is, without optical aid), those easily seen with binoculars, and those requiring a telescope to be appreciated. **Note, all of the objects (except single stars) will appear more impressive when viewed through a telescope or very large binoculars.** They are grouped in this way to highlight objects that can be seen using the optical equipment that may be available to the star gazer.

Tips for Observing the Night Sky

When observing the night sky, and in particular deep-sky objects such as star clusters, nebulae, and galaxies, it's always best to observe from a dark location. Avoid direct light from street lights and other sources. If possible observe from a dark location away from the light pollution that surrounds many of today's large cities.

You will see more stars after your eyes adapt to the darkness—usually about 10 to 20 minutes after you go outside. Also, if you need to use a torch to view the sky map, cover the light bulb with red cellophane. This will preserve your dark vision.

Finally, even though the Moon is one of the most stunning objects to view through a telescope, its light is so bright that it brightens the sky and makes many of the fainter objects very difficult to see. So try to observe the evening sky on moonless nights around either New Moon or Last Quarter.

Astronomical Glossary

Conjunction – An alignment of two celestial bodies such that they present the least angular separation as viewed from Earth.

Constellation – A defined area of the sky containing a star pattern.

Diffuse Nebula – A cloud of gas illuminated by nearby stars.

Double Star – Two stars that appear close to each other in the sky; either linked by gravity so that they orbit each other (binary star) or lying at different distances from Earth (optical double). Apparent separation of stars is given in seconds of arc (").

Ecliptic – The path of the Sun's center on the celestial sphere as seen from Earth.

Elongation – The angular separation of two celestial bodies. For Mercury and Venus the greatest elongation occurs when they are at their most angular distance from the Sun as viewed from Earth.

Galaxy – A mass of up to several billion stars held together by gravity.

Globular Star Cluster – A ball-shaped group of several thousand old stars.

Light Year (ly) – The distance a beam of light travels at 300,000 km/sec in one year.

Magnitude – The brightness of a celestial object as it appears in the sky.

Open Star Cluster – A group of tens or hundreds of relatively young stars.

Opposition – When a celestial body is opposite the Sun in the sky.

Planetary Nebula – The remnants of a shell of gas blown off by a star.

Universal Time (UT) – A time system used by astronomers. USA Eastern Standard Time (for example, New York) is 5 hours behind UT.

Variable Star – A star that changes brightness over a period of time.

NORTHERN HEMISPHERE JANUARY 2006 CELESTIAL OBJECTS Sky maps.com

Easily Seen with the Naked Eye

Capella	Aur	• The 6th brightest star. Appears yellowish in color. Spectroscopic binary. Dist=42 ly.
Sirius	CMa	• The brightest star in the sky. Also known as the "Dog Star". Dist=8.6 ly.
Procyon	CMi	• Greek name meaning "before the dog" - rises before Sirius (northern latitudes). Dist=11.4 ly.
δ Cephei	Cep	• Cepheid prototype. Mag varies between 3.5 & 4.4 over 5.366 days. Mag 6 companion.
Deneb	Cyg	• Brightest star in Cygnus. One of the greatest known supergiants. Dist=3,000 ly.
Castor	Gem	• Multiple star system with 6 components. 3 stars visible in telescope. Dist=52 ly.
Pollux	Gem	• With Castor, the twin sons of Leda in classical mythology. Dist=34 ly.
Vega	Lyr	• The 5th brightest star in the sky. A blue-white star. Dist=25.3 ly.
Rigel	Ori	• The brightest star in Orion. Blue supergiant star with mag 7 companion. Dist=770 ly.
Betelgeuse	Ori	• One of the largest red supergiant stars known. Diameter=300 times that of Sun. Dist=430 ly.
Algol	Per	• Famous eclipsing binary star. Magnitude varies between 2.1 & 3.4 over 2.867 days.
Pleiades	Tau	• The Seven Sisters. Spectacular cluster. Many more stars visible in binoculars. Dist=380 ly.
Hyades	Tau	• Large V-shaped star cluster. Binoculars reveal many more stars. Dist=151 ly.
Aldebaran	Tau	• Brightest star in Taurus. It is not associated with the Hyades star cluster. Dist=65 ly.
Polaris	UMi	• The North Pole Star. A telescope reveals an unrelated mag 8 companion star.

Easily Seen with Binoculars

M31	And	• The Andromeda Galaxy. Most distant object visible to naked eye. Dist=2.93 million ly.
M2	Aqr	• Resembles a fuzzy star in binoculars.
M38	Aur	• Stars appear arranged in "pi" or cross shape. Dist=4,300 ly.
M36	Aur	• About half size of M38. Located in rich Milky Way star field. Dist=4,100 ly.
M37	Aur	• Very fine star cluster. Discovered by Messier in 1764. Dist=4,400 ly.
M44	Cnc	• Praesepe or Beehive Cluster. Visible to the naked eye. Dist=577 ly.
M41	CMa	• First recorded observation by Aristotle in 325 BC as "cloudy spot". Dist=2,300 ly.
μ Cephei	Cep	• Herschel's Garnet Star. One of the reddest stars. Mag 3.4 to 5.1 over 730 days.
Mira	Cet	• Famous long period variable star. Mag varies between 3.0 & 10.1 over 332 days.
χ Cygni	Cyg	• Long period pulsating red giant. Magnitude varies between 3.3 & 14.2 over 407 days.
M39	Cyg	• May be visible to the naked eye under good conditions. Dist=900 ly.
v Draconis	Dra	• Wide pair of white stars. One of the finest binocular pairs in the sky. Dist=100 ly.
M35	Gem	• Fine open cluster located near foot of the twin Castor. Dist=2,800 ly.
γ Leporis	Lep	• Visible with binoculars. Gold & white stars. Mags 3.6 & 6.2. Dist=30 ly. Sep=96.3".
R Lyrae	Lyr	• Semi-regular variable. Magnitude varies between 3.9 & 5.0 over 46.0 days.
2232	Mon	• A large scattered star cluster of 20 stars. Dist=1,300 ly.
2244	Mon	• Surrounded by the rather faint Rosette Nebula. Dist=5,540 ly.
M50	Mon	• Visible with binoculars. Telescope reveals individual stars. Dist=3,000 ly.
Cr 69	Ori	• Lambda Orionis Cluster. Dist=1,630 ly.
M42	Ori	• The Great Orion Nebula. Spectacular bright nebula. Best with telescope. Dist=1,500 light years.
M15	Peg	• Only globular known to contain a planetary nebula (Mag 14, d=1"). Dist=30,000 ly.
Double Cluster	Per	• Double Cluster in Perseus. NGC 869 & 884. Excellent in binoculars. Dist=7,300 ly.
253	Scl	• Fine, large, cigar-shaped galaxy. Requires dark sky. Member of Sculptor Group.
Mizar & Alcor	UMa	• Good eyesight or binoculars reveals 2 stars. Not a binary. Mizar has a mag 4 companion.

Telescopic Objects

γ Andromedae	And	• Attractive double star. Bright orange star with mag 5 blue companion. Sep=9.8".
γ Arietis	Ari	• Impressive looking double blue-white star. Visible in a small telescope. Sep=7.8".
M67	Cnc	• Contains 500+ stars mag 10 & fainter. One of the oldest clusters. Dist=2,350 ly.
η Cassiopeiae	Cas	• Yellow star mag 3.4 & orange star mag 7.5. Dist=19 ly. Orbit=480 years. Sep=12".
61 Cygni	Cyg	• Attractive double star. Mags 5.2 & 6.1 orange dwarfs. Dist=11.4 ly. Sep=28.4".
γ Delphini	Del	• Appear yellow & white. Mags 4.3 & 5.2. Dist=100 ly. Struve 2725 double in same field.
θ Eridani	Eri	• Striking blue-white double star. Mags 3.2 & 4.3. Visible in a small telescope. Sep=8.2".
β Monocerotis	Mon	• Triple star. Mags 4.6, 5.0 & 5.4. Requires telescope to view arc-shape. Sep=7.3".
2264	Mon	• Christmas Tree Cluster. Associated with the Cone Nebula. Dist=2,450 ly.
σ Orionis	Ori	• Superb multiple star. 2 mag 7 stars one side, mag 9 star on other. Struve 761 triple in field.
M1	Tau	• Crab Nebula. Remnant from supernova which was visible in 1054. Dist=6,500 ly.
M33	Tri	• Fine face-on spiral galaxy. Requires a large aperture telescope. Dist=2.3 million ly.
M81	UMa	• Beautiful spiral galaxy visible with binoculars. Easy to see in a telescope.
M82	UMa	• Close to M81 but much fainter and smaller.