

About the Celestial Objects

Listed on this page are several of the brighter, more interesting celestial objects visible in the evening sky this month (refer to the monthly sky map). The objects are grouped into three categories. Those that can be easily seen with the naked eye (that is, without optical aid), those easily seen with binoculars, and those requiring a telescope to be appreciated. Note, all of the objects (except single stars) will appear more impressive when viewed through a telescope or very large binoculars. They are grouped in this way to highlight objects that can be seen using the optical equipment that may be available to the star gazer.

Tips for Observing the Night Sky

When observing the night sky, and in particular deep-sky objects such as star clusters, nebulae, and galaxies, it's always best to observe from a dark location. Avoid direct light from street lights and other sources. If possible observe from a dark location away from the light pollution that surrounds many of today's large cities.

You will see more stars after your eyes adapt to the darkness—usually about 10 to 20 minutes after you go outside. Also, if you need to use a torch to view the sky map, cover the light bulb with red cellophane. This will preserve your dark vision.

Finally, even though the Moon is one of the most stunning objects to view through a telescope, its light is so bright that it brightens the sky and makes many of the fainter objects very difficult to see. So try to observe the evening sky on moonless nights around either New Moon or Last Quarter.

Astronomical Glossary

Conjunction – An alignment of two celestial bodies such that they present the least angular separation as viewed from Earth.

Constellation – A defined area of the sky containing a star pattern.

Diffuse Nebula – A cloud of gas illuminated by nearby stars.

Double Star – Two stars that appear close to each other in the sky; either linked by gravity so that they orbit each other (binary star) or lying at different distances from Earth (optical double). Apparent separation of stars is given in seconds of arc (").

Ecliptic – The path of the Sun's center on the celestial sphere as seen from Earth.

Elongation – The angular separation of two celestial bodies. For Mercury and Venus the greatest elongation occurs when they are at their most angular distance from the Sun as viewed from Earth.

Galaxy – A mass of up to several billion stars held together by gravity.

Globular Star Cluster – A ball-shaped group of several thousand old stars.

Light Year (ly) – The distance a beam of light travels at 300,000 km/sec in one year.

Magnitude – The brightness of a celestial object as it appears in the sky.

Open Star Cluster – A group of tens or hundreds of relatively young stars.

Opposition – When a celestial body is opposite the Sun in the sky.

Planetary Nebula – The remnants of a shell of gas blown off by a star.

Universal Time (UT) - A time system used by astronomers. USA Eastern Standard Time (for example, New York) is 5 hours behind UT.

Variable Star – A star that changes brightness over a period of time.

NORTHERN HEMISPHERE MARCH 2006

OBJE(

ESTIAL

Easily Seen with the Naked Eye

•		•
Capella	Aur	• The 6th brightest star. Appears yellowish in color. Spectroscopic binary. Dist=42 ly.
Arcturus	Boo	 Orange, giant K star. Name means "bear watcher". Dist=37 ly.
Sirius	CMa	 The brightest star in the sky. Also known as the "Dog Star". Dist=8.6 ly.
Procyon	CMi	• Greek name meaning "before the dog" - rises before Sirius (northern latitudes). Dist=11.4 ly.
δ Cephei	Cep	Cepheid prototype. Mag varies between 3.5 & 4.4 over 5.366 days. Mag 6 companion.
Castor	Gem	 Multiple star system with 6 components. 3 stars visible in telescope. Dist=52 ly.
Pollux	Gem	• With Castor, the twin sons of Leda in classical mythology. Dist=34 ly.
Regulus	Leo	 Brightest star in Leo. A blue-white star with at least 1 companion. Dist=77 ly.
Rigel	0ri	• The brightest star in Orion. Blue supergiant star with mag 7 companion. Dist=770 ly.
Betelgeuse	0ri	• One of the largest red supergiant stars known. Diameter=300 times that of Sun. Dist=430 ly.
Algol	Per	 Famous eclipsing binary star. Magnitude varies between 2.1 & 3.4 over 2.867 days.
Pleiades	Tau	The Seven Sisters. Spectacular cluster. Many more stars visible in binoculars. Dist=380 ly.
Hyades	Tau	Large V-shaped star cluster. Binoculars reveal many more stars. Dist=151 ly.
Aldebaran	Tau	• Brightest star in Taurus. It is not associated with the Hyades star cluster. Dist=65 ly.
Polaris	UMi	• The North Pole Star. A telescope reveals an unrelated mag 8 companion star.

Easily Seen with Binoculars

, -			
M31	And	0	The Andromeda Galaxy. Most distant object visible to naked eye. Dist=2.93 million ly.
M38	Aur	0	Stars appear arranged in "pi" or cross shape. Dist=4,300 ly.
M36	Aur	0	About half size of M38. Located in rich Milky Way star field. Dist=4,100 ly.
M37	Aur	0	Very fine star cluster. Discovered by Messier in 1764. Dist=4,400 ly.
M44	Cnc	0	Praesepe or Beehive Cluster. Visible to the naked eye. Dist=577 ly.
M3	CVn	\oplus	Easy to find in binoculars. Might be glimpsed with the naked eye.
M41	CMa	0	First recorded observation by Aristotle in 325 BC as "cloudy spot". Dist=2,300 ly.
Mel 111	Com	0	Coma Berenices. 80 mag 5-6 stars in 5 deg. Dist=288 ly. Age=400 million years.
M35	Gem	0	Fine open cluster located near foot of the twin Castor. Dist=2,800 ly.
M48	Hya	0	12+ stars in 7x binoculars. Triangular asterism near centre. Dist=1,990 ly.
γ Leporis	Lep	•	Visible with binoculars. Gold & white stars. Mags 3.6 & 6.2. Dist=30 ly. Sep=96.3".
2232	Mon	0	A large scattered star cluster of 20 stars. Dist=1,300 ly.
2244	Mon	0	Surrounded by the rather faint Rosette Nebula. Dist=5,540 ly.
M50	Mon	0	Visible with binoculars. Telescope reveals individual stars. Dist=3,000 ly.
Cr 69	0ri	0	Lambda Orionis Cluster. Dist=1,630 ly.
M42	0ri		The Great Orion Nebula. Spectacular bright nebula. Best with telescope. Dist=1,500 light years.
Double Cluster	Per	0	Double Cluster in Perseus. NGC 869 & 884. Excellent in binoculars. Dist=7,300 ly.
M47	Pup	0	Bright star cluster. 15+ stars in 7x binoculars. Dist=1,500 ly.
M46	Pup	0	Dist=5.400 ly. Contains planetary NGC 2438 (Mag 11, d=65") - not associated.

Dist=5,400 ly. Contains planetary NGC 2438 (Mag 11, d=65") - not associated.

Mizar & Alcor UMa • Good eyesight or binoculars reveals 2 stars. Not a binary. Mizar has a mag 4 companion.

Telescopic Objects

rereacoh	Telescopic Objects						
γ Andromedae γ Arietis	And Ari	•	Attractive double star. Bright orange star with mag 5 blue companion. Sep=9.8". Impressive looking double blue-white star. Visible in a small telescope. Sep=7.8".				
ε Boötis	Воо	•					
M67	Cnc	0	Contains 500+ stars mag 10 & fainter. One of the oldest clusters. Dist=2,350 lv.				
M94	CVn	0					
M51	CVn	0	Whirlpool Galaxy. First recognised to have spiral structure. Dist=25 million ly.				
η Cassiopeiae	Cas	•	Yellow star mag 3.4 & orange star mag 7.5. Dist=19 ly. Orbit=480 years. Sep=12".				
M64	Com	0	Black-Eye Galaxy. Discovered by J.E. Bode in 1775 - "a small, nebulous star".				
3242	Hya		Ghost of Jupiter. Bright blue disk. Mag 11 central star. Dist=2,600 ly.				
γ Leonis	Leo	•	Superb pair of golden-yellow giant stars. Mags 2.2 & 3.5. Orbit=600 years. Sep=4.4".				
β Monocerotis	Mon	•	Triple star. Mags 4.6, 5.0 & 5.4. Requires telescope to view arc-shape. Sep=7.3".				
2264	Mon	0	Christmas Tree Cluster. Associated with the Cone Nebula. Dist=2,450 ly.				
σ Orionis	0ri	•	Superb multiple star. 2 mag 7 stars one side, mag 9 star on other. Struve 761 triple in field.				
k Puppis	Pup	•	Telescope easily shows two blue-white stars of almost equal brightness. Sep=9.9".				
M1	Tau		Crab Nebula. Remnant from supernova which was visible in 1054. Dist=6,500 ly.				
M81	UMa	0	Beautiful spiral galaxy visible with binoculars. Easy to see in a telescope.				
M82	UMa	0	Close to M81 but much fainter and smaller.				
γ Virginis	Vir	•	Superb pair of mag 3.5 yellow-white stars. Orbit=169 years. At their closest in 2005.				

